

2

LEVEL II

f

SOME GENERALIZATIONS OF KANTOROVICH INEQUALITY*

C. G. Khatri
Gujarat University

and

C. Radhakrishna Rao**
University of Pittsburgh

January 1981

Technical Report No. 81-1

DTIC
ELECTE
S APR 23 1981 D
F

Institute for Statistics and Applications
Department of Mathematics and Statistics
University of Pittsburgh
Pittsburgh, PA. 15260

**The work of the author is sponsored by the Air Force
Office of Scientific Research under Contract F49620-
79-0161. Reproduction in whole or in part is permitted
for any purpose of the United States Government.

*Based on a lecture delivered at the Conference on
Mathematical Statistics and Probability held at
New Delhi in December 1980 in honor of Professor
C. R. Rao on the occasion of his sixtieth birthday.

DTIC FILE COPY

SK

81 4 22 057

Approved for public release;
distribution unlimited.

**Best
Available
Copy**

REPORT DOCUMENTATION PAGE REPORT NUMBER AFOSR-TR-81-8385		LEAD INSTRUCTIONS BEFORE COMPLETING FORM 1. IF THIS IS A FINAL REPORT, CHECK BOX 1. 2. IF THIS IS A PRELIMINARY REPORT, CHECK BOX 2.	
TITLE (and Subtitle) Some Generalizations of Kantorovich Inequality		TYPE OF REPORT & PERIOD COVERED INTERIM Rept.	
AUTHOR C. G. Khatri and C. R. Rao <i>Radhakrishna</i>		U.P. Stat. Rept. 81-1 ✓ CONTRACT OR GRANT NUMBER F49620-79-C-0161	
PERFORMING ORGANIZATION NAME AND ADDRESS University of Pittsburgh Department of Mathematics & Statistics Pittsburgh, PA. 15260		PROGRAM ELEMENT, PROJECT, TASK, AND WORK UNIT NUMBER 2301/457 01102F	
CONTROLLING OFFICE NAME AND ADDRESS AFOSR/NM Bolling AFB. DC 20332		REPORT DATE January 1981	
MONITORING AGENCY NAME AND ADDRESS TR-81-1		NUMBER OF PAGES 18 4320	
DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited			
DISTRIBUTION STATEMENT (of the abstract entered in block 20, if different from Report) 			
SUPPLEMENTARY NOTES 			
KEY WORDS (Continue on reverse side if necessary; use block numbers) Kantorovich inequality, Least squares estimation			
ABSTRACT (Continue on reverse side if necessary; use block numbers) <p>Kantorovich gave an upper bound to $(x'Vx) (x'V^{-1}x)$ where x is an n-vector of unit length and V is an $n \times n$ positive definite matrix. Bloomfield, Watson and Knott found the bound to $x'VXX'V^{-1}x$, and Khatri and Rao to the trace and determinant of $X'VYY'V^{-1}X$ where X and Y are $n \times k$ matrices such that $X'X = Y'Y = I$. In the present paper we establish bounds for traces and determinants of $X'VYY'V^{-1}X$ and $X'BY'V'CX$ and $X'BY'Y'CX$</p>			

DD FORM 1473 JAN 73

UNCLASSIFIED

JCO

411385

81 4 22 057

~~UNCLASSIFIED~~

when X and Γ are matrices of different orders. A review of previous results on generalizations of Kantorovich inequality and a number of new results of independent interest are also given.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

SOME GENERALIZATIONS OF KANTOROVICH INEQUALITY

C. G. Khatri
Gujarat University
Ahmedabad, India

C. Radhakrishna Rao
University of Pittsburgh
Pittsburgh, PA. 15260

Summary: Kantorovich gave an upper bound to $(x'Vx)(x'V^{-1}x)$ where x is an n -vector of unit length and V is an $n \times n$ positive definite matrix. Bloomfield, Watson and Knott found the bound to $|X'VXX'V^{-1}X|$, and Khatri and Rao to the trace and determinant of $X'VYY'V^{-1}X$ where X and Y are $n \times k$ matrices such that $X'X = Y'Y = I$. In the present paper we establish bounds for traces and determinants of $X'VYY'V^{-1}X$ and $X'BY Y'CX$ when X and Y are matrices of different orders. A review of previous results on generalizations of Kantorovich inequality and a number of new results of independent interest are also given.

Key Words: Kantorovich inequality, Least squares estimation.

Accession For-	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)
NOTICE OF TRANSMITTAL TO DDC
This technical report has been reviewed and is
approved for public release IAW AFR 190-12 (7b).
Distribution is unlimited.
A. D. BLOSE
Technical Information Officer

1. INTRODUCTION

Let V be an $n \times n$ positive definite matrix with eigen values $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n > 0$, and define $\omega_i = (\lambda_i + \lambda_{n+i-1})^2 / 4 \lambda_i \lambda_{n+i-1}$, $i = 1, \dots, \alpha(\leq n/2)$. Kantorovich (1948) established the inequality

$$1 \leq (x' V x) (x' V^{-1} x) / (x' x)^2 \leq \omega_1 \quad (1.1)$$

for all non-null vectors x . A natural generalization of (1.1) is to compare the matrices $A = (X'X)^{-1} X' V X (X'X)^{-1}$ and $B = (X' V^{-1} X)^{-1}$ when X varies over $n \times k$ matrices of rank k . Let $\theta_1, \dots, \theta_k$ be the roots of $|A - \theta B| = 0$, i.e., the eigen values of A with respect to B . It is easy to establish that $\theta_i \geq 1$ for all i . Bloomfield and Watson (1975) and Knott (1975) showed that

$$|AB^{-1}| = \theta_1 \dots \theta_k \leq \prod_{i=1}^{\min(k, n-k)} \omega_i \quad (1.2)$$

while Khatri and Rao (1980) established that

$$\begin{aligned} \text{tr}(AB^{-1}) = \theta_1 + \dots + \theta_k &\leq \omega_1 + \dots + \omega_k \quad \text{if } n \geq 2k \\ &\leq \omega_1 + \dots + \omega_{n-k} + (2k-n), \quad \text{if } n < 2k \end{aligned} \quad (1.3)$$

where $\text{tr} C$ represents the trace of matrix C . It may be noted that $\sum \theta_i = \text{tr } P_X V P_X V^{-1}$ where P_X is the projection operator on the column space of X . Bloomfield and Watson (1975) gave another inequality

$$\text{tr } P_X V (I - P_X) V \leq \frac{1}{4} \sum_{i=1}^k (\lambda_i - \lambda_{n-i+1})^2 \quad (1.4)$$

when $n \geq 2k$. The inequalities (1.2) - (1.4) are useful in comparing the efficiencies of simple least squares estimators with the minimum variance unbiased estimators of parameters in the Gauss-Markoff model (see Khatri and Rao, 1980).

Khatri (1978, 1980) considered the matrix $(I - BA^{-1})$ which arises in a different context and proved the following results. Let $\underline{i} = (i_1, i_2, \dots, i_n)$ be a permutation of $(1, 2, \dots, n)$ and P denote the class of all permutations of $(1, 2, \dots, n)$. Further, let

$$\xi_{\underline{i}(\alpha)} = (\lambda_{i_\alpha} - \lambda_{i_{n-\alpha+1}})^2 / (\lambda_{i_\alpha} + \lambda_{i_{n-\alpha+1}})^2 \quad (1.5)$$

for $\alpha = 1, 2, \dots, k$ with $n \geq 2k$. Then

$$|I - BA^{-1}| \leq \sup_{\underline{i} \in P} \prod_{\alpha=1}^{\min(k, n-k)} \xi_{\underline{i}(\alpha)} = \prod_{j=1}^{\min(k, n-k)} \frac{(\lambda_j - \lambda_{n-j+1})^2}{(\lambda_j + \lambda_{n-j+1})^2}, \quad (1.6)$$

$$\text{tr } (I - BA^{-1}) \leq \sup_{\underline{i} \in P} \sum_{\alpha=1}^k \xi_{\underline{i}(\alpha)} \quad \text{if } n \geq 2k, \quad (1.7)$$

$$\text{tr } (I - BA^{-1})^{-1} \geq \inf_{\underline{i} \in P} \sum_{\alpha=1}^k \xi_{\underline{i}(\alpha)}^{-1} \quad \text{if } n \geq 2k. \quad (1.8)$$

Khatri (1978) further showed that if

$$C = X' V Y (Y' V Y)^{-1} Y' V X (X' V X)^{-1} \quad (1.9)$$

where Y is an $n \times s$, ($s \leq n-k$), matrix such that $X'Y = 0$,
then

$$|C| \leq |Q|, \text{tr } C \leq \text{tr } Q, \text{ and } \text{tr } C^{-1} \geq \text{tr } Q^{-1} \quad (1.10)$$

where $Q = I - BA^{-1}$. Eaton (1976) established the results (1.6) and (1.10) when $k=s=1$.

Strang (1960) generalized the Kantorovich inequality (1.1) in the form

$$[(x'Ax)(y'A^{-1}x) / (x'x)(y'y)] \leq \omega_1 \quad (1.11)$$

for all non-null vectors x and y where A is an $n \times n$ nonsingular matrix with singular values $\delta \geq \delta_2 \geq \dots \geq \delta_n > 0$ and

$$\omega_1 = (\delta_1 + \delta_{n-1+1})^2 / 4 \delta_1 \delta_{n-1+1}. \quad (1.12)$$

Greub and Rheinboldt (1959) proved that

$$\frac{(x'G^2x)(x'H^2x)}{(x'GHx)^2} \leq \frac{(\lambda_1\mu_1 + \lambda_n\mu_n)^2}{4\lambda_1\lambda_n\mu_1\mu_n} \quad (1.13)$$

for all non-null vectors x , when G and H are positive definite commuting matrices with eigen values $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n > 0$ and $\mu_1 \geq \mu_2 \geq \dots \geq \mu_n > 0$ respectively. The result (1.13) can be proved using (1.11).

A natural generalization of the expression in (1.11) is of the form

$$g(X,Y) = |X' A P_Y A^{-1} X| / |X' X| \quad (1.14)$$

or

$$f(X,Y) = \text{tr} (P_X A P_Y A^{-1}) \quad (1.15)$$

where X and Y are $(n \times k)$ and $(n \times s)$ matrices of ranks k and s respectively with $s \geq k$, and A is an $n \times n$ non-singular matrix with singular values $\delta_1 \geq \delta_2 \geq \dots \geq \delta_n > 0$. When $s = k$, Khatri and Rao (1980) established the inequalities

$$g(X,Y) \leq \prod_{i=1}^{\min(k, n-k)} \omega_i \quad (1.16)$$

$$\begin{aligned} f(X,Y) &\leq \sum_{i=1}^k \omega_i \quad \text{if } n \geq 2k \\ &\leq \sum_{i=1}^{n-k} \omega_i + 2k - n \quad \text{if } n < 2k \end{aligned} \quad (1.17)$$

where ω_i are as defined in (1.12). In the next section we establish the bounds for (1.14) and (1.15) when s is not necessarily equal to k .

We also consider determinants and traces of matrices of the type

$$(X' G^2 Y) (Y' G H Y)^{-1} (Y' H^2 X) (X' G H X)^{-1} \quad (1.18)$$

which are natural extensions of the expression in (1.13) and establish bounds under very general conditions on X , Y , G and H .

2. THE MAIN THEOREMS

In all the theorems stated in this section, X and Y stand for $n \times k$ and $n \times s$ matrices of ranks k and s respectively with $s \geq k$ and P_Z stands for the projection operator on the column space of matrix Z .

Theorem 1. Let $\delta_1 \geq \delta_2 \geq \dots \geq \delta_n > 0$ be the singular values of an $n \times n$ nonsingular matrix A , and

$$\omega_i = \frac{(\delta_i + \delta_{n-i+1})^2}{4\delta_i \delta_{n-i+1}}, \quad i = 1, \dots, m(\leq n/2). \quad (2.1)$$

Then

$$\frac{|X' A P_Y A^{-1} X|}{|X' X|} \leq \prod_{i=1}^{\min(k, n-s)} \omega_i \quad (2.2)$$

and

$$\begin{aligned} \text{tr}(P_X A P_Y A^{-1}) &\leq \sum_{i=1}^k \omega_i \quad \text{if } n \geq s+k \\ &\leq \sum_{i=1}^{n-s} \omega_i + (s+k-n) \quad \text{if } n < s+k \end{aligned} \quad (2.3)$$

Note 1. If X_1 and Y_1 are $n \times (n-k)$ and $n \times (n-s)$ matrices which are orthogonal complements of X and Y respectively, then $(n-k) \geq (n-s)$ and

$$\frac{|X' A P_Y A^{-1} X|}{|X' X|} = \frac{|Y_1' A^{-1} P_{X_1} A Y_1|}{|Y_1' Y_1|} \quad (2.4)$$

and

$$\text{tr} (P_X A P_Y A^{-1}) = (s+k-n) + \text{tr} (P_{Y_1} A^{-1} P_{X_1} A). \quad (2.5)$$

The results (2.4) and (2.5) show that we need only consider the case $s \geq k$ and $n \geq s+k$ in proving Theorem 1. If $n < s+k$, then $n \geq (n-s) + (n-k)$ in which case we consider the expressions on the right-hand sides of (2.4) and (2.5) and apply the same proof.

Note 2. If $A = P D_\delta Q'$ is the singular value decomposition of A , then we can write the left-hand side expressions of (2.2) and (2.3) as

$$|X'_* D_\delta Y_* Y'_* D_\delta^{-1} X_*| \text{ and } \text{tr} (X'_* D_\delta Y Y'_* D_\delta^{-1} X_*) \quad (2.6)$$

choosing $X_* = P'X(X'X)^{-\frac{1}{2}}$ and $Y_* = Q'Y(Y'Y)^{-\frac{1}{2}}$ so that $X'_* X_* = I_k$ and $Y'_* Y_* = I_s$, and $D_\delta = \text{Diag}(\delta_1, \dots, \delta_n)$ (i.e., a diagonal matrix with $\delta_1, \dots, \delta_n$ as diagonal elements). In view of (2.6), we need only prove Theorem 1 with the restrictions $X'X = I_k$, $Y'Y = I_s$ and A is a diagonal matrix.

Theorem 2. Let X and Y be $n \times k$ and $m \times s$ matrices of ranks k and s respectively with $k \leq s$, and B and C be $n \times m$ and $m \times n$ matrices such that $C = B^+$ (the Moore-Penrose inverse of B , see Rao, 1973, p. 26). Further, let $\delta_1 \geq \delta_2 \geq \dots \geq \delta_t > 0$ be the nonzero singular values of B and $t \geq s$. Then

$$\frac{|X' B P_Y C X|}{|X' X|} \leq \prod_{i=1}^{\min(k, t-s)} \omega_i \quad (2.7)$$

and

$$\begin{aligned} \text{tr}(P_X B P_Y C) &\leq \sum \omega_i \quad \text{if } t \geq s+k \\ &\leq s+k-t + \sum_{i=1}^{t-s} \omega_i \quad \text{if } t < s+k \end{aligned} \quad (2.8)$$

where $\omega_i = (\delta_i + \delta_{t-i+1})^2 / 4 \delta_i \delta_{t-i+1}$.

Theorem 3. Let V and W be $n \times n$ and $m \times m$ non-negative definite matrices, and X and Y be $n \times k$ and $m \times s$ matrices such that $X' V X$ and $Y' W Y$ are positive definite. Further, let B and C be $n \times m$ and $m \times n$ matrices such that

$$(a) \quad t = R(B) = R(C) \leq s$$

$$(b) \quad \rho(W) = \rho\left(\frac{B}{W}\right) = \rho(C:V), \quad \rho(V) = \rho(B:V) = \rho\left(\frac{C}{V}\right)$$

where $\rho(A)$ stands for the rank of matrix A .

$$(c) \quad BW^+ C \quad \text{and} \quad CV^+ B \quad \text{are symmetric of rank } t \\ \text{and} \quad BW^+ CV^+ B = B.$$

If $\delta_1^2 \geq \delta_2^2 \geq \dots \geq \delta_t^2 > 0$ are the nonzero eigen values of $BW^+ B' V^+$ and $\omega_i = (\delta_i + \delta_{t-i+1})^2 / 4 \delta_i \delta_{t-i+1}$, then

$$\frac{|X' B Y (Y' W Y)^{-1} Y' C X|}{|X' V X|} \leq \prod_{i=1}^{\min(k, t-s)} \omega_i \quad (2.9)$$

and

$$\begin{aligned} \text{tr}[(X' V X)^{-1} X' B Y (Y' W Y)^{-1} Y' C X] &\leq \sum_{i=1}^k \omega_i, \quad \text{if } t \geq s+k \\ &\leq (s+k-t) + \sum_{i=1}^{t-s} \omega_i, \quad \text{if } t < s+k. \end{aligned} \quad (2.10)$$

Theorem 4. Let S and R be $n \times m$ and $m \times n$ matrices such that $t = \rho(S) = \rho(R) = \rho(SR)$ with $t \geq s \geq k$, and SR and RS are symmetric, nonnegative definite and idempotent matrices. Further let $\delta_1 \geq \delta_2 \geq \dots \geq \delta_t > 0$ be the nonzero singular values of S and $\omega_i = (\delta_i + \delta_{t-i+1})^2 / 4\delta_i \delta_{t-i+1}$. Then

$$\frac{|X' S Y (Y' R S Y)^{-1} Y' R X|}{|X' S R X|} \leq \prod_{i=1}^{\min(k, t-s)} \omega_i \quad (2.11)$$

and

$$\begin{aligned} \text{tr}[(X' S R X)^{-1} X' S Y (Y' R S Y)^{-1} Y' R X] \\ \leq \sum_{i=1}^k \omega_i \quad \text{if } t \geq s + k \\ \leq (s + k - t) + \sum_{i=1}^{t-s} \omega_i \quad \text{if } t < s + k. \end{aligned} \quad (2.12)$$

Proof. Theorem 4 follows from Theorem 3 by choosing $V = SR$, $W = RS$, $B = S$ and $C = R$.

Theorem 5. Let G and H be symmetric and commuting matrices such that GH is nonnegative definite and $\rho(G) = \rho(H) = \rho(GH) = t \geq s \geq k$. Further let $\delta_1 \geq \delta_2 \geq \dots \geq \delta_t > 0$ be the nonzero eigen values of $H^{-1}G$ where H^{-1} is any g -inverse of H , and $\omega_i = (\delta_i + \delta_{t-i+1})^2 / 4\delta_i \delta_{t-i+1}$. Then

$$\frac{|X' G^2 Y (Y' G H Y)^{-1} Y' H^2 X|}{|X' G H X|} \leq \prod_{i=1}^{\min(k, t-s)} \omega_i \quad (2.13)$$

and

$$\begin{aligned} & \text{tr}[(X'GHX)^{-1}X'G^2Y(Y'GHY)^{-1}Y'H^2X] \\ & \leq \sum_{i=1}^k \omega_i \quad \text{if } t \geq k+s \\ & \leq (k+s-t) + \sum_{i=1}^{t-s} \omega_i \quad \text{if } t < k+s. \quad (2.14) \end{aligned}$$

Proof. Theorem 5 follows from Theorem 3 by choosing $V = W = GH$, $B = G^2$ and $C = H^2$.

Theorem 6. Let V and W be positive definite, and B and C be two matrices such that $\rho(B) = \rho(C) = t (\geq s)$, $BW^{-1}C$ and $CV^{-1}B$ are symmetric and $BW^{-1}CV^{-1}B = B$. Further let $\delta_1 \geq \delta_2 \geq \dots \geq \delta_t > 0$ be the nonzero singular values of $V_1^{-1}B(W_1')^{-1}$ where $W = W_1W_1'$ and $V = V_1V_1'$. Then the inequalities (2.9) and (2.10) hold.

3. PROOFS OF MAIN THEOREMS

Proof of Theorem 1. The proof depends on a number of lemmas which are also of independent interest. As observed in Notes 1 and 2 following the statement of Theorem 1, we can take X and Y such that $X'X = I_k$, $Y'Y = I_s$, $n \geq s+k$ and A as $D_\delta = \text{Diag}(\delta_1, \dots, \delta_n)$ with all δ_i positive.

Lemma 1. Let V be an n.n.d. matrix of order n with non-zero eigen values $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_s > 0$ and X be an $n \times k$ matrix of rank k ($\leq s$). Then

$$\sup_X \frac{|X'VX|}{|X'X|} = \prod_{i=1}^k \lambda_i \quad \text{and} \quad \sup_X \text{tr}(P_X V) = \sum_{i=1}^k \lambda_i \quad (3.1)$$

Proof. The result (3.1) is an immediate consequence of the Poincaré Separation Theorem (see Rao, 1979, p. 364) which states that

$$\mu_i \leq \lambda_i, \quad i = 1, \dots, k (\leq s) \quad (3.2)$$

where $\mu_i, i = 1, 2, \dots$ are the roots of the determinantal equation

$$|X'VX - \mu X'X| = 0 \quad (3.3)$$

and the equality in (3.2) is attained for a suitably chosen X .

Lemma 2. Let X and Y be $n \times k$ and $n \times s$ matrices such that $X'X = I_k$, $Y'Y = I_s$ and $s \geq k$. Let D_δ be a positive definite diagonal matrix and $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_s > 0$ be the eigen values of $(Y'D_\delta Y)(Y'D_\delta^{-1} Y)$. Then

$$|X'D_\delta Y Y'D_\delta^{-1} X| \leq \left(\prod_{i=1}^k \lambda_i \right)^{\frac{1}{2}} |X'D_\delta X X'D_\delta^{-1} X|^{\frac{1}{2}} \quad (3.4)$$

and

$$\text{tr}(P_X D_\delta P_Y D_\delta^{-1}) \leq \left(\sum_{i=1}^k \lambda_i \right)^{\frac{1}{2}} (\text{tr} P_X D_\delta P_X D_\delta^{-1})^{\frac{1}{2}} \quad (3.5)$$

Proof of (3.4). Consider

$$|X'D_{\delta} Y Y' D_{\delta}^{-1} X|^2 = |(DX)' B (DX)|$$

where $D^2 = D_{\delta}$ and

$$B = D Y Y' D_{\delta}^{-1} X X' D_{\delta}^{-1} Y Y' D$$

which is an n.n.d. matrix of rank k . If $\alpha_1, \dots, \alpha_k$ are the nonzero eigen values of B , then

$$\left(\prod_{i=1}^k \alpha_i \right) = |X'D_{\delta}^{-1} Y Y' D_{\delta}^{-1} X|. \quad (3.6)$$

Hence using the result (3.1) of Lemma 1 with X as DX and V as B we have

$$|X'D_{\delta} Y Y' D_{\delta}^{-1} X|^2 \leq |X'D_{\delta} X| \cdot |X'D_{\delta}^{-1} Y Y' D_{\delta} Y Y' D_{\delta}^{-1} X|. \quad (3.7)$$

Now,

$$|X'D_{\delta}^{-1} Y Y' D_{\delta} Y Y' D_{\delta}^{-1} X| = |(D^{-1}X)' C (D^{-1}X)| \quad (3.8)$$

where $C = D^{-1} Y Y' D_{\delta} Y Y' D^{-1}$ is an n.n.d. matrix of rank s and its eigen values are $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_s > 0$. Then, again applying the result (3.1) to the expression on the right hand side of (3.8), we have from (3.7)

$$|X'D_{\delta} Y Y' D_{\delta}^{-1} X|^2 \leq |X'D_{\delta} X| \cdot |X'D^{-1} X| \left(\prod_{i=1}^k \lambda_i \right)$$

which proves (3.4).

Proof of (3.5). Consider the singular value decompositions

$$D^{-1} X X' D = T_1 D_1 T_2' \quad \text{and} \quad D Y Y' D^{-1} = T_3 D_2 T_4'$$

where T_1, T_2, T_3 and T_4 are orthogonal matrices and D_1 and D_2 are diagonal matrices such that $D_1 = \text{Diag}(\beta_1, \dots, \beta_k, 0, \dots)$ and $D_2 = \text{Diag}(\gamma_1, \dots, \gamma_s, 0, \dots)$, where $\beta_1 \geq \dots \geq \beta_k > 0$ and $\gamma_1 \geq \dots \geq \gamma_s > 0$. Then using a theorem of von Neumann (see equation (2.11) in Rao, 1979),

$$\text{tr}(P_X D_\delta P_Y D_\delta^{-1}) = \text{tr}(D_1 T_2' T_3 D_2 T_4' T_1) \leq \text{tr} D_1 D_2 \quad (3.9)$$

and the equality holds iff

$$T_2' T_3 = \begin{pmatrix} I_k & 0 \\ 0 & \Delta_1 \end{pmatrix} \quad \text{and} \quad T_4' T_1 = \begin{pmatrix} I_k & 0 \\ 0 & \Delta_2 \end{pmatrix}$$

where Δ_1 and Δ_2 are arbitrary orthogonal matrices.

Now,

$$\text{tr} D_1 D_2 = \sum_{i=1}^k \gamma_i \beta_i \leq \left(\sum_{i=1}^k \gamma_i^2 \right)^{\frac{1}{2}} \left(\sum_{i=1}^k \beta_i^2 \right)^{\frac{1}{2}}, \quad (3.10)$$

$$\sum_{i=1}^k \beta_i^2 = \text{tr}(P_X D_\delta P_X D_\delta^{-1})$$

and $\lambda_i = \gamma_i^2, i = 1, \dots, s$ are the nonzero eigen values of $P_Y D_\delta P_Y D_\delta^{-1}$. Hence using (3.10) in (3.9) we get

$$\text{tr}(P_X D_\delta P_Y D_\delta^{-1}) \leq (\text{tr} P_X D_\delta P_X D_\delta^{-1})^{\frac{1}{2}} \left(\sum_{i=1}^k \lambda_i \right)^{\frac{1}{2}}$$

which proves (3.5). Thus Lemma 2 is established.

Lemma 3. Let $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_s > 0$ be the eigen values of $Y' D_\delta Y Y' D_\delta^{-1} Y$ where Y is an $n \times s$ matrix such that $Y'Y = I_s$ and $\text{diag } D_\delta = (\delta_1, \dots, \delta_n)$ with all δ_i positive, and $\omega_i = (\delta_i + \delta_{n-i+1})^2 / 4\delta_i \delta_{n-i+1}$ for $i = 1, \dots, k (\leq s)$. Then

$$\prod_{i=1}^k \lambda_i \leq \prod_{i=1}^k \omega_i \quad \text{and} \quad \sum_{i=1}^k \lambda_i \leq \sum_{i=1}^k \omega_i. \quad (3.11)$$

Proof. Let

$$\begin{aligned} \phi(B) &= |B'Y' D_\delta Y B| / |B'(Y' D_\delta^{-1} Y)^{-1} B| \\ \psi(B) &= \text{tr}[B'Y' D_\delta Y B (B'(Y' D_\delta^{-1} Y)^{-1} B)^{-1}] \end{aligned} \quad (3.12)$$

where B is an $s \times k$ matrix with $\rho(B) = k \leq s$. Then applying (3.1)

$$\sup_B \phi(B) = \prod_{i=1}^k \lambda_i \quad \text{and} \quad \sup_B \psi(B) \leq \sum_{i=1}^k \lambda_i. \quad (3.13)$$

Next we observe that

$$B'(Y' D_\delta^{-1} Y)^{-1} B - B'B(B'Y' D_\delta^{-1} Y B)^{-1} B'B \quad (3.14)$$

is non-negative definite. (see example 33 on p. 77 of Rao, 1973). Then, substituting the second expression in (3.14) for the first in (3.12), we get

$$\phi(B) \leq |(YB)' D_{\delta}(YB)| \quad |(YB)' D_{\delta}^{-1}(YB)| / |(YB)'(YB)|^2$$

$$\Psi(B) \leq \text{tr}\{(YB)' D_{\delta}(YB) | (YB)' YB |^{-1} (YB)' D_{\delta}^{-1}(YB) | (YB)'(YB) |^{-1}\}. \quad (3.15)$$

Now, writing $YB = L$ which is of rank k and applying the inequalities (1.2) and (1.3) to the right hand sides of (3.15) we have

$$\phi(B) \leq \prod_{i=1}^k \omega_i \quad \text{and} \quad \Psi(B) \leq \sum_{i=1}^k \omega_i$$

which in conjunction with (3.13) proves (3.11), since ω_i are independent of B .

Combining the results (3.4) and (3.11) we get

$$\begin{aligned} |X' D_{\delta} Y Y' D_{\delta}^{-1} X| &\leq \left(\prod_{i=1}^k \omega_i \right)^{\frac{1}{2}} |X' D_{\delta} X X' D_{\delta}^{-1} X|^{\frac{1}{2}} \\ \text{tr}(X' D_{\delta} Y Y' D_{\delta}^{-1} X) &\leq \left(\sum_{i=1}^k \omega_i \right)^{\frac{1}{2}} (\text{tr} X' D_{\delta} X X' D_{\delta}^{-1} X)^{\frac{1}{2}}. \end{aligned} \quad (3.16)$$

A further application of (1.2) and (1.3) to the right hand side expressions of (3.16) proves Theorem 1.

Proof of Theorem 2. Consider the singular value decomposition, $B = \Delta_1' D_{\delta} \Delta_2'$ where $\Delta_i' \Delta_i = I_t$ for $i=1,2$ and $D_{\delta} = \text{Diag}(\delta_1, \dots, \delta_t)$ with all δ_i positive. Then $C = B^+ = \Delta_2 D_{\delta}^{-1} \Delta_1'$. Let $X_0 = \Delta_1' X$ and $Y_0 = \Delta_2' Y$ so that $X'X - X_0' X_0$ and $Y'Y - Y_0' Y_0$ are n.n.d. matrices and so also

$$P_{Y_0} = Y_0 (Y'Y)^{-1} Y_0' \quad \text{and} \quad P_{X_0} = X_0 (X'X)^{-1} X_0'.$$

Then it is easily seen that

$$\frac{|X' B P_Y C X|^2}{|X' X|^2} \leq \frac{|X'_O D_\delta P_{Y_O} D_\delta^{-1} X_O|^2}{|X'_O X_O|^2} \quad (3.17)$$

and

$$\text{tr}(P_X B P_Y C) \leq \text{tr}(P_{X_O} D_\delta P_{Y_O} D_\delta^{-1}). \quad (3.18)$$

Now using Theorem 1 on the right hand side expressions of (3.17) and (3.18), we get the results (2.7) and (2.8) of Theorem 2.

Proof of Theorem 3. Let V and W be of ranks n_1 and m_1 respectively and write $V = V_1 V_1'$ and $W = W_1 W_1'$ where V_1 and W_1 are $n \times n_1$ and $m \times m_1$ matrices of ranks n_1 and m_1 respectively. Let

$$B_1 = (V_1' V_1)^{-1} V_1' B W_1 (W_1' W_1)^{-1} \text{ and } C_1 = (W_1' W_1)^{-1} W_1' C V_1 (V_1' V_1)^{-1}.$$

Then, under the given conditions, it is easy to verify that $C_1 = B_1^+$ and taking $X_O = V_1' X$ and $Y_O = W_1' Y$,

$$X' B Y (Y' W Y)^{-1} Y' C X = X'_O B_1 Y_O (Y'_O Y_O)^{-1} Y'_O C_1 X_O \quad (3.19)$$

and $X' V X = X'_O X_O$. Now applying Theorem 2 to the right-hand side expression of (3.19), we get the results of Theorem 3.

REFERENCES

- [1] Bloomfield, P. and Watson, G. S. (1975). The inefficiency of least squares. Biometrika 62 121-128.
- [2] Eaton, M. L. (1976). A maximization problem and its applications to canonical correlation. J. Multivar. Anal., 6 422-425.
- [3] Greub, W. and Rheinboldt, W. (1959). On a generalization of an inequality of L. V. Kantorovich. Proc. Amer. Math. Soc., 10 407-415.
- [4] Kantorovich, L. V. (1948). Functional analysis and applied mathematics. Uspehi Mat. Nauk., 3 89-135; (also translated from the Russian by C. D. Benster, Nat. Bur. Standards Report No. 1509, March 7, 1952).
- [5] Khatri, C. G. (1978). Some optimization problems with applications to canonical correlations and sphericity tests. J. Multivar. Anal., 8 453-467. Corrections. (1980). J. Multivar. Anal., 10.
- [6] Khatri, C. G. and Rao, C. R. (1980). Some extensions of Kantorovich inequality and statistical applications. To appear in J. Multivar. Anal., 10.
- [7] Knott, M. (1975). On the minimum efficiency of least squares. Biometrika 62 129-132.
- [8] Rao, C. R. (1973). Linear Statistical Inference and its Applications. 2nd edition Wiley, New York.
- [9] Rao, C. R. (1979). Separation theorems for singular values of matrices and their applications in multivariate analysis. J. Multivar. Anal., 9 362-377.
- [10] Strang, W. G. (1960). On the Kantorovich inequality. Proc. Amer. Math. Soc., 11 468.